


arXiv:hep-th/9804194v1 29 Apr 1998

Black Holes in the Framework of the Four-Dimensional Effective Theory of Heterotic Superstrings at Low Energies

Panagiota Kanti¹

*Physics Department, Division of Theoretical Physics,
University of Ioannina, Greece*



A Dissertation in Candidacy for the Degree of Doctor of Philosophy

February 1998

¹Email : pkanti@cc.uoi.gr . A full postscript version of the dissertation in Greek (197 pages) can be retrieved from the address : <http://artemis.sci.uoi.gr/~pkanti/phd.ps.gz>

Summary

Superstring Theory is the most promising unification scheme of the four forces of nature. On the other hand, black holes are the first and most exciting solutions of Einstein's equations and their properties are defined by various theorems of the Theory of General Relativity, mainly by the “No-hair” theorem. The Low Energy Effective Superstring Theory plays the role of the connecting link between these two theories and, at the same time, the role of a generalised gravitational theory. In the framework of this effective theory, we search for new black hole solutions. By extending existing treatments of black hole solutions in string gravity, we compute the analytical expressions of all the scalar fields of the effective theory, that is of the dilaton, the axions and the modulus field, outside the horizon of a Kerr-Newman black hole and in $\mathcal{O}(\alpha')$ [1]. In the presence of higher curvature gravitational terms, such as the quadratic Gauss-Bonnet term, we demonstrate analytically the evasion of the existing “no-hair” theorems including the novel one by Bekenstein. In addition, we determine via numerical integration the existence of a new family of black hole solutions, the so called Dilatonic Black Holes, which are dressed with classical dilaton hair [2]. In the framework of the more general Einstein-Dilaton-Gauss-Bonnet-Yang-Mills theory, another family of solutions, the Coloured Black Holes, are determined. These solutions are characterized by non-trivial dilaton and Yang-Mills hair for the particular case of $SU(2)$ gauge fields [3]. The thermodynamical properties of the black hole solutions are also discussed. The hair, however, found in all of the above cases is of “secondary type”, in the sense that it owes its existence to the primary gravitational or electromagnetic field. Finally, the linear stability of Dilatonic Black Holes under small spacetime-dependent perturbations is exhibited through a semi-analytic method which makes use of the Fubini-Sturm's theorem [4]. This result is extremely important in that it constitutes a linearly stable example of a black hole that bypasses the existing “no-hair” theorems.

Publications

- [1] “Classical Moduli $\mathcal{O}(\alpha')$ Hair”, P. Kanti and K. Tamvakis, Phys. Rev. **D 52** (1995) 3506-3511.
- [2] “Dilatonic Black Holes in Higher Curvature String Gravity”, P. Kanti, N.E. Mavromatos, J. Rizos, K. Tamvakis and E. Winstanley, Phys. Rev. **D 54** (1996) 5049-5058.
- [3] “Coloured Black Holes in Higher Curvature String Gravity”, P. Kanti and K. Tamvakis, Phys. Lett. **B 392** (1997) 30-38.
- [4] “Dilatonic Black Holes in Higher Curvature String Gravity II : Linear Stability”, P. Kanti, N.E. Mavromatos, J. Rizos, K. Tamvakis and E. Winstanley, Phys. Rev. **D 57** (to appear in the 15th May 1998 issue).

CONTENTS

Preface	1
1. Introduction to String Theory	3
1.1 Introduction	3
1.2 Bosonic String	4
1.3 Superstring	16
1.4 Heterotic Superstring and Compactification	29
1.5 Effective String Theory at Low Energies	36
1.6 Four-Dimensional Effective Theory and Moduli Fields	45
2. Introduction to Black Holes	49
2.1 Introduction	49
2.2 The Schwarzschild Metric.....	52
2.3 The Kerr-Newman Metric	60
2.4 Killing Vectors and Surface Gravity	64
2.5 Thermodynamics of Black Holes	66
2.6 The “No-hair” Theorem	73
3. “Scalar Hair” in Kerr-Newman Black Hole Background	81
3.1 Introduction	81
3.2 Action Functional and Equations of Motion of the Theory	86
3.3 Solution at Zero Order in α'	88
3.4 Solution at First Order in α'	90
Appendix 3.A Derivation of the Equations of Motion	97
Appendix 3.B The Kerr Metric	103
4. Black Holes with “Dilaton Hair” in the Presence of Higher Curvature Gravitational Terms	105
4.1 Introduction	105
4.2 Action Functional and Equations of Motion of the Theory	106
4.3 The Novel “No-hair” Theorem	111

4.4 Numerical Analysis	115
4.4.1 Determination of Black Hole Solutions	115
4.4.2 Determination of the Independent Parameters of the Black Hole Solutions at Infinity	119
4.4.3 Determination of Additional Solutions	122
Appendix 4.A The Coefficients d , d_1 and d_2	127
5. “Coloured” Black Holes in the Presence of Higher Curvature Gravitational Terms	129
5.1 Introduction	129
5.2 Action Functional and Equations of Motion of the Theory	130
5.3 Asymptotic Form of the Solutions at Infinity and the Event Horizon	133
5.4 Thermodynamical Study of Black Hole Solutions	137
5.5 Numerical Analysis	142
5.5.1 Determination of Black Hole Solutions	143
5.5.2 Determination of “Naked Singularities”	150
6. Stability Analysis of Black Holes with “Dilaton “Hair”	153
6.1 Introduction	153
6.2 Equations of Motion of the Time-Dependent Theory	154
6.3 Linear Stability Analysis	157
6.4 Asymptotic Behaviour of Perturbation $\delta\phi$	164
6.5 Application of the Fubini-Sturm’s Theorem	167
6.6 Stability of the “Coloured” Black Holes	172
7. Conclusions	177
Bibliography	185